

**College of the Holy Cross, Fall Semester, 2021**  
**Math 302** (Professor Hwang), Meeting 02  
**Space Paths, Arc Length**

The definitions of velocity, speed, acceleration, and regularity make sense for paths in  $\mathbf{R}^n$ . We will mostly consider paths in space ( $\mathbf{R}^3$ ).

If our path  $\alpha$  is regular on some interval containing  $t_0$ , we define its *arc length function* to be the integral of the speed,

$$s(t) = \int_{t_0}^t |\alpha'(u)| du.$$

(If  $t_0$  is unspecified, assume it is 0.) The *arc length* over  $[a, b]$  is  $s(b) - s(a)$ , the definite integral of the speed.

**Exercise 02.1:** Set up (but *do not attempt to evaluate* the arc length functions for the indicated paths.

(a)  $\alpha(t) = (at, bt^2)$ ;      (b)  $\alpha(t) = (a \cos t, b \sin t)$ ;      (c)  $\alpha(t) = (a \cosh t, b \sinh t)$ .

**Exercise 02.2:** Let  $r > 0$  and  $k$  be real. Calculate the arc length function of the *helix*

$$\alpha(t) = (r \cos t, r \sin t, kt),$$

and give an arc length parametrization.

**Exercise 02.3:** Let  $r_i > 0$  and  $k_i$  be real. Calculate the arc length function of the path

$$\alpha(t) = (r_1 \cos(k_1 t), r_1 \sin(k_1 t), r_2 \cos(k_2 t), r_2 \sin(k_2 t)).$$

**Exercise 02.4:** Calculate the arc length function of the parabola  $\alpha(t) = (t, \frac{1}{2}t^2)$ . Hint:  $\int \sqrt{1+t^2} dt = \frac{1}{2}[t\sqrt{1+t^2} + \ln(t + \sqrt{1+t^2})]$ .

**Exercise 02.5:** Each part refer to the *catenary*  $\alpha(t) = (t, \cosh t)$ .

(a) Calculate the arc length function.

(b) Suppose  $s = \sinh t = \frac{1}{2}(e^t - e^{-t})$ . Solve for  $t$  as a function of  $s$ . Hint: Multiply by  $e^t$  and move everything to one side. Then use the quadratic formula to solve for  $e^t$ .

(c) Give an arc length parametrization of the catenary. Hint:  $\cosh^2 u = 1 + \sinh^2 u$  for all  $u$ , so  $\cosh(\sinh^{-1} s) = \dots$ ?

If time permits, here are some exercises on polar graphs.

**Exercise 02.6:** Suppose  $I$  is an interval of real numbers, and that  $f$  is a differentiable, real-valued function on  $I$ . The *polar graph*  $r = f(\theta)$  may be parametrized by setting  $r = f(t)$  and  $\theta = t$  in polar coordinates:

$$\alpha(t) = (r \cos \theta, r \sin \theta) = f(t)(\cos t, \sin t).$$

Calculate the velocity and speed of  $\gamma$ . Under what conditions on  $f$  is the path  $\gamma$  regular? Hint: You can crunch out the speed by brute force, but if you use vectors and dot products you may save yourself some work!

**Exercise 02.7:** Let  $R$  be a positive real number, and let  $\alpha$  be the parametrization of the polar graph  $r = 2R \cos \theta$ .

- (a) Find a Cartesian equation for the image of  $\gamma$ . Describe the polar graph geometrically. Suggestion: Multiply both sides by  $r$ , then convert to Cartesian.
- (b) Use your formulas from the preceding question to calculate the velocity and speed of  $\gamma$ . Is anything noteworthy?

**Exercise 02.8:** Calculate the arc length of the polar graph  $r = 1 + \cos \theta$ . Hint:  $\sqrt{2 + 2 \cos \theta} = 2 \cos \frac{\theta}{2}$  by the double-angle formula for cosine.