

**College of the Holy Cross, Fall Semester, 2005**  
**Math 131, Practice Midterm 3 Solutions**

1. Match each parametric curve with its graph. (Each graph shows the square  $-1 \leq x \leq 1$ ,  $-1 \leq y \leq 1$ .)

(a)  $x(t) = -1 + 2t$ ,  $y(t) = 1 - 2t$ ,  $0 \leq t \leq 1$        I    II    III    IV    V

**Explanation.** This is a line with slope  $-2/2 = -1$ .

(b)  $x(t) = \cos(2t)$ ,  $y(t) = \sin(2t)$ ,  $0 \leq t \leq \pi$        I    II    III    IV    V


**Explanation.** Points  $(x, y)$  on the curve satisfy  $x^2 + y^2 = 1$ , so they lie on the unit circle.

(c)  $x(t) = \cos t$ ,  $y(t) = \frac{1}{2} \sin t$ ,  $0 \leq t \leq 2\pi$        I    II    III    IV    V


**Explanation.** This is the standard parametrization of the unit circle, but with the  $y$ -coordinate scaled by a factor of  $1/2$ , so it must be an ellipse.

(d)  $x(t) = 1 - 2t$ ,  $y(t) = -t$ ,  $0 \leq t \leq 1$        I    II    III    IV    V


**Explanation.** This is a line with slope  $-1/-2 = 1/2$ .




I




II



III



IV



V

2. Compute the indicated limits. Show all work for full credit.

(a)  $\lim_{x \rightarrow 1} \frac{\cos(\frac{\pi}{2}x)}{x-1}$

**Answer.** Since  $\cos(\pi/2) = 0$ , the numerator and denominator both approach zero. By L'Hopital's Rule,

$$\lim_{x \rightarrow 1} \frac{\cos(\frac{\pi}{2}x)}{x-1} = \lim_{x \rightarrow 1} \frac{-\frac{\pi}{2} \sin(\frac{\pi}{2}x)}{1} = -\frac{\pi}{2}$$

Limit =

$-\frac{\pi}{2}$

(b)  $\lim_{x \rightarrow 0} \frac{x^2 + 1}{x^2 + 3}$

**Answer.**

$$\lim_{x \rightarrow 0} \frac{x^2 + 1}{x^2 + 3} = \frac{\lim_{x \rightarrow 0} x^2 + 1}{\lim_{x \rightarrow 0} x^2 + 3} = \frac{1}{3}$$

Limit =

$\frac{1}{3}$
---------------

(c)  $\lim_{x \rightarrow \infty} \frac{x \ln x}{x^{1.01}}$  **Answer.** First simplify.

$$\lim_{x \rightarrow \infty} \frac{x \ln x}{x^{1.01}} = \lim_{x \rightarrow \infty} \frac{\ln x}{x^{0.01}}$$

The numerator and denominator both approach  $\infty$ , so by L'Hopital's Rule,

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x^{0.01}} = \lim_{x \rightarrow \infty} \frac{\frac{1}{x}}{0.01x^{-0.99}} = \lim_{x \rightarrow \infty} \frac{1}{0.01x^{0.01}} = 0$$

Limit =

0
---

3. Each part refers to the function  $f(x) = 2x^3 - 6x^2$ .

(a) [10 points] Find and classify (local min/max, or neither) the critical points of  $f$ .

**Answer.** The derivative of  $f$  is  $f'(x) = 6x^2 - 12x = 6x(x - 2)$ , so the critical points of  $f$  are  $x = 0$  and  $x = 2$ .

To determine the type of each critical point, apply either the first or second derivative test.

First Derivative Test. Since  $f'(-1) = 18$ ,  $f'(1) = -6$  and  $f'(3) = 18$ , the sign of  $f'$  changes from  $+$  to  $-$  at  $x = 0$ , so  $f$  has a local maximum at  $x = 0$ , and the sign of  $f'$  changes from  $-$  to  $+$  at  $x = 2$ , so  $f$  has a local minimum at  $x = 2$ .

Second Derivative Test. The second derivative of  $f$  is  $f''(x) = 12x - 12$ . Since  $f''(0) = -12$  is negative,  $f$  has a local maximum at  $x = 0$ , and since  $f''(2) = 12$  is positive,  $f$  has a local minimum at  $x = 2$ .

(b) [5 points] Find the maximum and minimum values of  $f(x)$  if  $-\frac{3}{2} \leq x \leq \frac{5}{2}$ .

**Answer.** To find the global maximum and minimum, evaluate  $f$  at each critical point and at each endpoint:


$$\begin{aligned} f(0) &= 0 \\ f(2) &= -8 \\ f\left(-\frac{3}{2}\right) &= -20.25 \\ f\left(\frac{5}{2}\right) &= -6.25 \end{aligned}$$

Minimum value: - 20.25

Maximum value: 0

- (c) [5 points] In the grid provided, sketch the graph  $y = f(x)$  for  $-1 \leq x \leq 3$ . (Note that axis labels are provided.) For full credit, clearly indicate the critical and inflection point(s) in this interval, and label each such point with *both coordinates*.

**Answer.**  $f''(x) = 12x - 12 = 12(x - 1)$ , so  $f''$  changes sign at  $x = 1$ . Therefore  $f$  has an inflection point at  $(1, -4)$ . The critical points have coordinates  $(0, 0)$  and  $(2, -8)$ .



4. [15 points] Crusader Movie Rentals finds that they can rent 160 movies per night at \$1 per movie. For every dollar that the rental fee increases, 40 fewer movies are rented. What price should be charged to maximize the revenue (total rental income)?

**Answer.** Let  $p$  be the price in dollars per movie, and let  $n$  be the number of movies that can be rented at that price. The revenue is then  $R = np$ . Let  $x$  be the increase in the price from 1 dollar. Then  $p = 1 + x$ , and  $n = 160 - 40x$ , so

$$R(x) = (1 + x)(160 - 40x) = 160 + 120x - 40x^2.$$

The domain of  $R$  is  $-1 \leq x \leq 4$ . Since

$$R'(x) = 120 - 80x,$$

the critical point of  $R$  is  $x = \frac{3}{2} = 1.5$ . Since  $R(-1) = R(4) = 0$  and  $R(1.5) = 250$ , the revenue is maximized when  $x = 1.5$ , i.e. when  $p = 2.5$ .

Revenue maximized when  $p =$  \$2.50

---

5. [15 points] Find the equation of the line tangent to the curve  $xy - 2x - 3y + 1 = 0$  at the point  $(-2, 1)$ .

**Answer.** Use implicit differentiation:

$$x \frac{dy}{dx} + 1 \cdot y - 2 - 3 \frac{dy}{dx} = 0$$

Now solve for  $\frac{dy}{dx}$ :

$$\frac{dy}{dx} = \frac{2 - y}{x - 3}$$


When  $x = -2$  and  $y = 1$ ,

$$\frac{dy}{dx} = \frac{1}{-5} = -\frac{1}{5}.$$

So the slope of the tangent line is  $m = -\frac{1}{5}$ . Applying the point-slope formula gives  $y - 1 = -\frac{1}{5}(x + 2)$ .

Equation of tangent line at  $(-2, 1)$ :  $y - 1 = -\frac{1}{5}(x + 2)$  or  $y = -\frac{1}{5}x + \frac{3}{5}$

6. A boat is drawn into a dock by a rope over a small pulley. The pulley is five feet higher than the bow of the boat (see figure). Let  $\ell$  be the length of rope,  $x$  the distance from the boat to the dock.



- (a) [5 points] Find an equation relating  $\ell$  and  $x$ , and determine  $x$  when  $\ell = 13$ .

**Answer.** Use the Pythagorean Theorem.

Equation relating  $\ell$  and  $x$ :  $x^2 + 25 = \ell^2$

When  $\ell = 13$ ,  $x =$  12

- (b) [10 points] Suppose the rope is drawn in at 3 ft/sec. How fast is the boat moving when the length of the rope is 13 feet?

**Answer.** We are given  $\frac{d\ell}{dt} = -3$  and want to find  $\frac{dx}{dt}$  when  $\ell = 13$ . Differentiate the equation  $x^2 + 25 = \ell^2$  with respect to  $t$ :

$$2x \frac{dx}{dt} = 2\ell \frac{d\ell}{dt}$$

When  $\ell = 13$ ,  $x = 12$ , so

$$2(12) \frac{dx}{dt} = 2(13)(-3)$$

$$\frac{dx}{dt} = -3.25$$

Speed = 3.25 ft/sec