

MATH 135 – Calculus 1  
Second Derivative and Concavity  
November 18, 2016

*Background*

We say  $f$  (or the graph  $y = f(x)$ ) is *concave up* on an interval if  $f'$  is increasing on that interval, and similarly,  $f$  or its graph is *concave down* if  $f'$  is decreasing on that interval. Combined with our results from last time, this says:

- If  $f''(x) > 0$  on an interval, then  $f$  or its graph is concave up on that interval
- If  $f''(x) < 0$  on an interval, then  $f$  or its graph is concave down on that interval
- A point  $(c, f(c))$  on the graph of  $f$  where the concavity changes is called a *point of inflection* of  $f$ .

The notion of concavity can also be used to state a second method for determining whether critical points are local maxima or local minima, called the Second Derivative Test:

**Theorem 1** (*Second Derivative Test*) *Let  $f$  be differentiable on some open interval containing a critical point  $c$ . In addition, assume  $f''(c)$  exists.*

- (a) *If  $f''(c) > 0$ , then  $f(c)$  is a local minimum*
- (b) *If  $f''(c) < 0$ , then  $f(c)$  is a local maximum*
- (c) *If  $f''(c) = 0$ , there is no conclusion.*

In the last case here,  $f$  could have either a local maximum or a local minimum, or neither, so no conclusion is possible. *Technical Comment:* In the other cases, the intuition is that  $f'$  should be increasing or decreasing on an interval containing  $c$  depending on the sign of  $f''(c) = (f')'(c)$ , so that (a) corresponds to a case where the graph is concave up at  $c$  and (b) corresponds to a case where the graph is concave down at  $c$ . This would follow, for instance, if we knew (in addition) that  $f''$  was continuous on some interval containing  $c$ . But the conclusion of the Theorem is valid even without that extra continuity hypothesis, as is shown in Exercise 67 in Section 4.4.

*Questions*

1. Consider the graph  $f(x) = x^3 - 3x^2 + 2x$  on the interval  $[-1, 3]$  from last time (the plot is also on the back of this sheet). Find the intervals where  $f$  is concave up and the intervals where  $f$  is concave down. How many points of inflection are there on this graph and where are they located?
2. Consider  $f(x) = x^2e^{-x}$ .
  - (a) Compute  $f'(x)$  and find all critical points.
  - (b) Determine the sign of  $f'(x)$  on each interval between successive critical points, and use that to classify the critical points as local maxima or local minima by the First Derivative Test.




Figure 1: Plot for question 1

- (c) Now compute  $f''(x)$  and check your answers in (b) by using the Second Derivative Test.
  - (d) Determine all points of inflection of  $f$ .
3. Repeat question 2 for  $f(x) = 2x^4 - 3x^2 + 2$ .