

MATH 135 – Calculus 1
The Squeeze Theorem and Trigonometric Limits
October 3, 2016

Background

In today's video, we saw an additional technique for evaluating limits called the "Squeeze Theorem" and the very important limit:

$$\lim_{\theta \rightarrow 0} \frac{\sin(\theta)}{\theta} = 1. \quad (1)$$

Recall that the Squeeze Theorem says: Assume $l(x) \leq f(x) \leq u(x)$ on some interval containing $x = c$ (except possibly at $x = c$) and $\lim_{x \rightarrow c} l(x) = \lim_{x \rightarrow c} u(x) = L$ (we might say " f is squeezed by l and u at $x = c$ " to describe this). Then $\lim_{x \rightarrow c} f(x)$ exists and equals L as well.

Questions

Do the following problems from Section 2.6 in our text:

- (1) Suppose the graphs $y = f(x)$, $y = l(x)$, $y = u(x)$ are as in the plot on the back. What can we say about $\lim_{x \rightarrow 1} f(x)$?
- (2) Determine $\lim_{x \rightarrow 0} f(x)$ given that $\cos(x) \leq f(x) \leq 1$ for all x .
- (3) State whether the given inequality provides sufficient information to determine $\lim_{x \rightarrow 1} f(x)$, and if so, find the limit. (Hint: Draw pictures!)
 - (a) $4x - 5 \leq f(x) \leq x^2$
 - (b) $2x - 1 \leq f(x) \leq x^2$
 - (c) $4x - x^2 \leq f(x) \leq x^2 + 2$

Evaluate the following limits using (1):

(4)
$$\lim_{t \rightarrow 0} \frac{\sin(t)}{8t}.$$

(5)
$$\lim_{t \rightarrow 0} \frac{\sin(8t)}{t}.$$

(Hint: For this one let $u = 8t$ and convert the t in the denominator to an equivalent expression in terms of u . Note that $t \rightarrow 0$ implies $u = 8t \rightarrow 0$ also.)

(6)
$$\lim_{t \rightarrow 0} \frac{\sin(3t)}{\sin(5t)}.$$




Figure 1: Plot for Question 1, $y = l(x), u(x)$ in blue; $y = f(x)$ in red.

(Hint: Rewrite as follows:

$$\frac{\sin(3t)}{\sin(5t)} = \frac{\frac{\sin(3t)}{t}}{\frac{\sin(5t)}{t}},$$

then proceed as in question (5).

(7)

$$\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t^2}$$