

MATH 133 – Calculus with Fundamentals 1  
Trigonometric Limits  
October 6, 2017

*Background*

In today's video, we saw an additional technique for evaluating limits called the "Squeeze Theorem" and the very important limit:

$$\lim_{\theta \rightarrow 0} \frac{\sin(\theta)}{\theta} = 1. \quad (1)$$

Recall that the Squeeze Theorem says: Assume  $l(x) \leq f(x) \leq u(x)$  on some interval containing  $x = c$  (except possibly at  $x = c$ ) and  $\lim_{x \rightarrow c} l(x) = \lim_{x \rightarrow c} u(x) = L$  (we might say " $f$  is squeezed by  $l$  and  $u$  at  $x = c$ " to describe this). Then  $\lim_{x \rightarrow c} f(x)$  exists and equals  $L$  as well.

*Questions*

Evaluate the following limits using (1):

(1)

$$\lim_{t \rightarrow 0} \frac{\sin(t)}{8t}.$$

(2)

$$\lim_{t \rightarrow 0} \frac{\sin(8t)}{t}.$$

(Hint: For this one let  $u = 8t$  and convert the  $t$  in the denominator to an equivalent expression in terms of  $u$ . Note that  $t \rightarrow 0$  implies  $u = 8t \rightarrow 0$  also.)

(3)

$$\lim_{t \rightarrow 0} \frac{\sin(3t)}{\sin(5t)}.$$

(Hint: Rewrite as follows:

$$\frac{\sin(3t)}{\sin(5t)} = \frac{\frac{\sin(3t)}{t}}{\frac{\sin(5t)}{t}},$$

then proceed as in question (2).