## MATH 133 – Calculus with Fundamentals 1 The Derivative Function October 18, 2017

## Background

Recall that last time, we introduced the *derivative* of a function f at x:

$$f'(x) = \lim_{h \to 0} \frac{f(x+h) - f(x)}{h},$$

provided the limit exists.

In today's video we introduced the idea of deriving formulas for f'(x) for a general x, and considering f' as a new function in its own right. We said:

- (1) If  $f(x) = x^n$  for any number n, then  $f'(x) = nx^{n-1}$ .
- (2) If f'(x) exists, then so does (kf)'(x) and (kf)'(x) = kf'(x).
- (3) If f'(x) and g'(x) both exist, then so does (f+g)'(x), and (f+g)'(x) = f'(x) + g'(x).

## Questions

- (1) Compute f'(x) for  $f(x) = x^{5/3}$  using point (1) in the Background above.
- (2) Compute f'(x) for  $f(x) = x^4 + 5x^2 + 7 + 2x^{-3}$  using points (1), (2), and (3) in the Background above.
- (3) Make up your own example function of the same type as the one in (2) and compute f'(x) using points (1), (2), and (3) in the Background.
- (4) What is f'(x) if  $f(x) = \pi^{\pi}$ ? (Be careful!)

The video for today gave the idea for showing point (1) in the Background when n is a positive integer, and stated that the same formula works for all n. Let's verify this in some special cases.

- (5) Verify the formula in point (1) for  $f(x) = x^4$ . That is, you want to show  $f'(x) = 4x^3$  using the definition of f'(x).
- (6) Verify the formula in point (1) for  $f(x) = \frac{1}{x^2} = x^{-2}$ . That is, you want to show  $f'(x) = -2x^{-3}$  using the definition of f'(x).
- (7) Why does point (2) in the background work? Explain in terms of the limit definition, and in terms of how the graphs y = f(x) and y = kf(x) are related.

Finally, it is important to realize that the limit in the definition of f'(x) can fail to exist at some of the x in the domain of f.

(9) Consider y=|x| (the absolute value function). Does

$$f'(0) = \lim_{h \to 0} \frac{|h| - 0}{h}$$

exist? Why or why not? Does this make sense in terms of the graph y = |x|?