


College of the Holy Cross, Spring Semester, 2018
MATH 134 Solutions for Practice Final Exam
May 3, 2018

I.

(A) Let $f(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 2 \\ x - 1 & \text{if } 2 < x \leq 4 \\ 11 - 2x & \text{if } 4 < x \leq 6 \end{cases}$ whose graph is shown here:

Solution:



Let $F(x) = \int_{-1}^x f(t) dt$, where $f(t)$ is the function from part (B). (There is a part of the graph to the left of $x = 0$ that you are not seeing. Complete the following table of values for $F(x)$):

Solution: The value $F(x)$ represents the signed area between the graph $y = f(x)$ and the x -axis. By the interval union property for integrals

$$\int_{-1}^x f(x) dx = \int_{-1}^0 f(x) dx + \int_0^x f(x) dx$$

and we can tell $\int_{-1}^0 f(x) dx = 3$ from the first entry in the table. Then to find the others we just compute areas and add that 3:

x	0	1	2	3	4	5	6
$F(x)$	3	4	5	6.5	9	11	11

(B) Compute the derivative of the function $g(x) = \int_0^{2x} \frac{\cos(t)}{t^2} dt$.

Solution: By the first part of the Fundamental Theorem of Calculus and the Chain Rule for derivatives:

$$g'(x) = \frac{\cos(2x)}{(2x)^2} \cdot 2 = \frac{\cos(2x)}{2x^2}.$$

II. Compute the following integrals.

(A) $\int \frac{x^4 - 3\pi^3 + \sqrt{x}}{x^{2/3}} dx$

Solution: Split into separate fractions, simplify and integrate:

$$\begin{aligned} \int \frac{x^4 - 3\pi^3 + \sqrt{x}}{x^{2/3}} dx &= \int x^{10/3} - 3\pi^3 x^{-2/3} + x^{-1/6} dx \\ &= \frac{3}{13} x^{13/3} - 9\pi^3 x^{1/3} + \frac{6}{5} x^{5/6} + C. \end{aligned}$$

(B) $\int x^3 e^{x^2} dx$ (integrate by parts)

Solution: The correct choice is $u = x^2$ and $dv = x e^{x^2} dx$ (note that you need the x with the exponential in order to integrate!) Hence $du = 2x dx$ and $v = \frac{1}{2} e^{x^2}$. Then by the parts formula

$$\int x^3 e^{x^2} dx = \frac{1}{2} x^2 e^{x^2} - \int x e^{x^2} dx = \frac{1}{2} x^2 e^{x^2} - \frac{1}{2} e^{x^2} + C.$$

(C) $\int \frac{\csc^2(5x) dx}{\cot(5x) + 7}$

Solution: This one can be handled by the u -substitution $u = \cot(5x) + 7$, for which $du = -5 \csc^2(5x) dx$ by the Chain Rule. Then

$$\int \frac{\csc^2(5x) dx}{\cot(5x) + 7} = \frac{-1}{5} \int u^{-1} du = \frac{-1}{5} \ln |u| + C = \frac{-1}{5} \ln |\cot(5x) + 7| + C.$$

(D) $\int_1^e x^5 \ln(x) dx$.

Solution: Integrate by parts with $u = \ln(x)$ and $dv = x^5 dx$. Then $du = \frac{1}{x} dx$ and $v = \frac{1}{6} x^6$ and by the integration by parts formula,

$$\int x^5 \ln(x) dx = \frac{x^6}{6} \ln(x) - \int \frac{1}{6} x^6 \cdot \frac{1}{x} dx = \frac{x^6}{6} \ln(x) - \frac{x^6}{36} + C.$$

For the definite integral, we apply the Fundamental Theorem to get

$$\int_1^e x^5 \ln(x) dx = \left. \frac{x^6}{6} \ln(x) - \frac{x^6}{36} \right|_1^e = \frac{e^6}{6} \ln(e) - \frac{e^6}{36} - \frac{1}{6} \ln(1) + \frac{1}{36} = \frac{5e^6 + 1}{36}.$$

$$(E) \int \frac{1}{\sqrt{16+x^2}} dx$$

Solution: This can be done by the trigonometric substitution $x = 4 \tan \theta$, so $dx = 4 \sec^2 \theta d\theta$, and $\sqrt{16+x^2} = \sqrt{16(1+\tan^2 \theta)} = 4 \sec \theta$. This simplifies to

$$\int \frac{1}{4 \sec \theta} \cdot 4 \sec^2 \theta d\theta = \int \sec \theta d\theta.$$

Now we use formula ST4 from our table, then convert back to x :

$$\begin{aligned} \int \sec \theta d\theta &= \ln |\sec \theta + \tan \theta| + C \\ &= \ln \left| \frac{\sqrt{x^2+16}}{4} + \frac{x}{4} \right| + C. \end{aligned}$$

Using properties of logarithms and incorporating $-\ln(4)$ with the constant, this can also be written in the form:

$$\ln |\sqrt{16+x^2} + x| + C.$$

$$(F) \int \frac{x}{(x^2+1)(x+1)} dx$$

Solution: Using partial fractions, we must solve for A, B, C to make:

$$\frac{1}{(x+1)(x^2+1)} = \frac{A}{x+1} + \frac{Bx+C}{x^2+1}.$$

Clearing denominators,

$$1 = A(x^2+1) + (Bx+C)(x+1),$$

so equating coefficients, $A+B=0$, $B+C=0$, and $A+C=1$. Solving simultaneously,

$$A = \frac{1}{2}, \quad B = \frac{-1}{2}, \quad C = \frac{1}{2}.$$

Then

$$\begin{aligned} \int \frac{1}{(x+1)(x^2+1)} dx &= \int \frac{\frac{1}{2}}{x+1} + \frac{\frac{-x}{2} + \frac{1}{2}}{x^2+1} dx \\ &= \frac{1}{2} \ln |x+1| + \frac{-1}{4} \ln(x^2+1) + \frac{1}{2} \tan^{-1} x + C. \end{aligned}$$

III.

(A) Use a midpoint Riemann sum with $n = 4$ to approximate the integral

$$\int_1^3 \sqrt{1+9x^4} dx.$$

Solution: The R_4 Riemann sum approximation is


$$\begin{aligned} \int_1^3 \sqrt{1+9x^4} dx &\doteq \left(\sqrt{1+9(1.5)^4} + \sqrt{1+9(2)^4} + \sqrt{1+9(2.5)^4} + \sqrt{1+9(3)^4} \right) (.5) \\ &\doteq 32.3302. \end{aligned}$$

Solution: The midpoint approximation is an *overestimate* because $\sqrt{1+9x^4}$ is increasing on $[1, 3]$.

IV. A region R in the plane is bounded by the graphs $y = 9 - x^2$, $y = 2x$, $x = 0$ and $x = 1$.

(B) Compute the area of the region R .

Solution: Here is a sketch of the region:



The area is given by the integral

$$A = \int_0^1 9 - x^2 - 2x dx = 9x - \frac{x^3}{3} - x^2 \Big|_0^1 = \frac{23}{3}.$$

(B) Compute the volume of the solid obtained by rotating R about the x -axis.

Solution: The cross-sections of the solid by planes perpendicular to the x -axis are washers with inner radius $r_{in} = 2x$ and outer radius $r_{out} = 9 - x^2$. So the volume is the integral of the area of the cross-section:

$$\begin{aligned} V &= \int_0^1 \pi(9 - x^2)^2 - \pi(2x)^2 dx = \pi \int_0^1 81 - 22x^2 + x^4 dx \\ &= \pi \left(81x - \frac{22x^3}{3} + \frac{x^5}{5} \Big|_0^1 \right) \\ &= \frac{1108\pi}{15}. \end{aligned}$$

(C) Set up the integral(s) to compute the volume of the solid obtained by rotating R about the y -axis. *You do not need to compute the value.*

Solution: The easiest way (by far) is to use cylindrical shells:

$$V = \int_0^1 2\pi x(9 - x^2 - 2x) dx$$

If you use slices, the cross-sections by planes perpendicular to the y -axis are all disks, but the function giving the radius of the disk is given by three different formulas depending on whether $0 \leq y \leq 2$, or $2 \leq y \leq 8$, or $8 \leq y \leq 9$.

$$V = \int_0^2 \pi \left(\frac{y}{2}\right)^2 dy + \int_2^8 \pi(1)^2 dy + \int_8^9 \pi(\sqrt{9-y})^2 dy.$$

V. The daily solar radiation x per square meter (in hundreds of calories) in Florida in October has a probability density function $f(x) = c(x-2)(6-x)$ if $2 \leq x \leq 6$, and zero otherwise. Find value of c and the probability that the daily solar radiation per square meter is greater than 4.

Solution: We must have

$$1 = \int_2^6 c(x-2)(6-x) dx = c \left(-\frac{x^3}{3} + 4x^2 - 12x \Big|_2^6 \right) = \frac{32c}{3}.$$

Therefore $c = \frac{3}{32}$. Then the probability that $x > 4$ is given by the integral

$$\bar{x} = \int_4^6 \frac{3}{32}(x-2)(6-x) dx = \int_4^6 \frac{3x}{4} - \frac{3x^2}{32} - \frac{9}{8} dx = \frac{1}{2}.$$

VI. An avian flu epidemic has broken out in Birdsburgh, a large city with total population 10 million. Let N be the number, in millions, of people who have been infected, as a function of time t in weeks. The Birdsburgh Public Health department proposes the model that the rate of change of N is proportional to the product of the number of infected people (N) and the number of people not yet infected.

- (A) (10) Write the proposed model above as a differential equation, calling the constant of proportionality k .

Solution: If N people have been infected (N in millions), then the number who have not been infected is $10 - N$ (millions). So the differential equation is

$$\frac{dN}{dt} = kN(10 - N).$$

(Note that this can be put into the form of a logistic equation:

$$\frac{dN}{dt} = (10k)N \left(1 - \frac{N}{10} \right);$$

the 10 plays the role of the carrying capacity.)

- (B) (5) The function $N(t) = 10/(1 + 9999e^{-t})$ should be a solution of your differential equation from part A. What is the value of k ?

Solution: For this N ,

$$\frac{dN}{dt} = \frac{-99990e^{-t}}{(1 + 9999e^{-t})^2} \quad (1)$$

and

$$\begin{aligned} N(10 - N) &= \frac{10}{1 + 9999e^{-t}} \cdot \left(10 - \frac{10}{1 + 9999e^{-t}}\right) \\ &= \frac{10 \cdot (-99990e^{-t})}{(1 + 9999e^{-t})^2} \end{aligned} \quad (2)$$

Hence comparing (1) and (2), we must have $k = \frac{1}{10} = .1$. (Note: This could also be done by recognizing that the differential equation can be rearranged to the form of a logistic equation with constant $10k$: $\frac{dN}{dt} = (10k)N \left(1 - \frac{N}{10}\right)$. The general solution of this is

$$N = \frac{10}{1 + ce^{-(10k)t}}$$

So if the given function is a solution we must have $c = 9999$ and $10k = 1$, so $k = .1$.)

- C) If the epidemic proceeds according the function given in part B, how many weeks will pass before the number of infected people reaches 1 million?

Solution: We must solve $1 = \frac{10}{1+9999e^{-t}}$. So $1 + 9999e^{-t} = 10$, and $t = -\ln(9/9999) = \ln(1111) \doteq 7.0$ weeks. Note that the units of N are millions of people so the left side of the equation is $N = 1$, not $N = 1000000$.

VII.

- (A) Find the sum of the finite geometric series with alternating signs:

$$3 - \frac{3}{2} + \frac{3}{2^2} - \frac{3}{2^3} + \cdots + \frac{3}{2^6}$$

What happens if you continue adding terms of the same form indefinitely? That is, what is the sum

$$\sum_{k=0}^{\infty} \frac{(-1)^k 3}{2^k}?$$

Solution: The geometric series has $c = 3$ and $a = \frac{-1}{2}$, so the sum is

$$\frac{3 \left(1 - \left(\frac{-1}{2}\right)^7\right)}{1 - \left(\frac{-1}{2}\right)} = \frac{3 \left(1 + \frac{1}{128}\right)}{\frac{3}{2}} = \frac{257}{128}.$$

The infinite sum is

$$\lim_{N \rightarrow \infty} \frac{3 \left(1 + \left(\frac{-1}{2}\right)^N\right)}{\frac{3}{2}} = 2.$$

- (B) What would be the monthly payment on a \$500,000 30-year fixed-rate mortgage at 3.5% annual interest.

Solution: The future value of the \$500000 is

$$FV = \$500000 \left(1 + \frac{.035}{12} \right)^{12 \cdot 30} = \$1,426,643.75$$

(rounding to the nearest cent). On the other hand a stream of monthly payments P extending over 30 years has future value

$$P \cdot \frac{\left(1 + \frac{.035}{12} \right)^{360} - 1}{1 + \frac{.035}{12} - 1} \doteq P \cdot 635.4128591$$

Setting the future values equal, we solve for P :

$$P = \frac{\$1,426,643.75}{635.4128591} \doteq \$2245.22$$

(You need to be quite well-off to be able to afford a loan this big!!)