

MATH 135 – Calculus 1  
The Derivative Chain Rule  
November 4, 2019

*Background*

Our next major derivative short-cut rule is *one of the most important*. This rule, called the *Chain Rule* allows us to differentiate functions that are built up by *composition*. Here's what it says: If  $g$  is differentiable at  $x$  and  $f$  is differentiable at  $g(x)$ , then the composition  $(f \circ g)(x) = f(g(x))$  is differentiable at  $x$  and

$$\frac{d}{dx}f(g(x)) = f'(g(x))g'(x).$$

In words: The derivative of the composition is *the derivative of the outside function (i.e.  $f'$ ), with  $g(x)$  “plugged in,” times the derivative of  $g$* . For example, the function

$$h(x) = \sqrt{x^2 + 4x + 9}$$

is the composition  $f(g(x))$  where  $f(x) = \sqrt{x}$  and  $g(x) = x^2 + 4x + 9$  is “plugged in.” The Chain Rule says the derivative of  $h$  will be given by computing  $f'(x) = \frac{1}{2\sqrt{x}}$  (do you see where that comes from?), plugging  $g$  into  $f'(x)$ , then multiplying by  $g'(x)$ :

$$h'(x) = \frac{1}{2\sqrt{x^2 + 4x + 9}} \cdot (2x + 4) = \frac{x + 2}{\sqrt{x^2 + 4x + 9}}.$$

Today's class will be devoted to understanding and practicing this rule. We'll continue and use this a different way next time.

*Questions*

For each function, identify an  $f(x)$  and  $g(x)$  such that the given function is the composition  $f(g(x))$ . Then apply the Chain Rule and compute the derivative:

(a)

$$h(x) = e^{3x+1}.$$

(b)

$$h(x) = \frac{1}{(x^4 + 5x^2 + 1)^{3/2}}.$$

(c)

$$h(x) = \sin(\cos(x) + x).$$

(d)

$$h(x) = (\tan(x) + 4x)^3.$$

(e) Sometimes we need to use the Chain Rule more than once (if the function we're looking at is “several composition layers deep” like

$$h(x) = \cos^2(4x^3 + 2) = (\cos(4x^3 + 2))^2.$$

Note that this is  $f(g(x))$  with  $f(x) = x^2$  and  $g(x) = \cos(4x^3 + 2)$ . But  $g(x)$  is also a composition, so you'll need to use the Chain Rule again to find  $g'(x)$ . With these hints, find  $h'(x)$ .