

MATH 135 – Calculus 1  
The Squeeze Theorem and Trigonometric Limits  
October 2, 2019

*Background*

In today's video, we saw an additional technique for evaluating limits called the "Squeeze Theorem" and the very important limit:

$$\lim_{\theta \rightarrow 0} \frac{\sin(\theta)}{\theta} = 1. \quad (1)$$

Recall that the Squeeze Theorem says: Assume  $l(x) \leq f(x) \leq u(x)$  on some interval containing  $x = c$  (except possibly at  $x = c$ ) and  $\lim_{x \rightarrow c} l(x) = \lim_{x \rightarrow c} u(x) = L$  (we might say " $f$  is squeezed by  $l$  and  $u$  at  $x = c$ " to describe this). Then  $\lim_{x \rightarrow c} f(x)$  exists and equals  $L$  as well.

*Questions*

Do the following problems from Section 2.6 in our text:

- (1) Suppose the graphs  $y = f(x)$ ,  $y = l(x)$ ,  $y = u(x)$  are as in the plot on the back. What can we say about  $\lim_{x \rightarrow 1} f(x)$ ?
- (2) Determine  $\lim_{x \rightarrow 0} f(x)$  given that  $\cos(x) \leq f(x) \leq 1$  for all  $x$ .
- (3) State whether the given inequality provides sufficient information to determine  $\lim_{x \rightarrow 1} f(x)$ , and if so, find the limit. (Hint: Draw pictures!)
  - (a)  $4x - 5 \leq f(x) \leq x^2$
  - (b)  $2x - 1 \leq f(x) \leq x^2$
  - (c)  $4x - x^2 \leq f(x) \leq x^2 + 2$

Evaluate the following limits using (1):

(4) 
$$\lim_{t \rightarrow 0} \frac{\sin(t)}{8t}.$$

(5) 
$$\lim_{t \rightarrow 0} \frac{\sin(8t)}{t}.$$

(Hint: For this one let  $u = 8t$  and convert the  $t$  in the denominator to an equivalent expression in terms of  $u$ . Note that  $t \rightarrow 0$  implies  $u = 8t \rightarrow 0$  also.)

(6) 
$$\lim_{t \rightarrow 0} \frac{\sin(3t)}{\sin(5t)}.$$




Figure 1: Plot for Question 1,  $y = l(x), u(x)$  in blue;  $y = f(x)$  in red.

(Hint: Rewrite as follows:

$$\frac{\sin(3t)}{\sin(5t)} = \frac{\frac{\sin(3t)}{t}}{\frac{\sin(5t)}{t}},$$

then proceed as in question (5).

(7)

$$\lim_{t \rightarrow 0} \frac{1 - \cos(t)}{t^2}$$