

ELEMENTS BOOK 2

Fundamentals of Geometric Algebra


Ὅροι.

α'. Πᾶν παραλληλόγραμμον ὀρθογώνιον περιέχεσθαι λέγεται ὑπὸ δύο τῶν τὴν ὀρθὴν γωνίαν περιεχουσῶν εὐθειῶν.

β'. Παντὸς δὲ παραλληλογράμμου χωρίου τῶν περὶ τὴν διάμετρον αὐτοῦ παραλληλογράμμων ἐν ὁποιοοῦν σὺν τοῖς δυοῖς παραπληρώμασι γνῶμων καλεῖσθω.

α'.

Ἐὰν ὡσι δύο εὐθεῖαι, τμηθῆ δὲ ἡ ἑτέρα αὐτῶν εἰς ὅσα δηποτοῦν τμήματα, τὸ περιεχόμενον ὀρθογώνιον ὑπὸ τῶν δύο εὐθειῶν ἴσον ἐστὶ τοῖς ὑπὸ τε τῆς ἀτμήτου καὶ ἐκάστου τῶν τμημάτων περιεχομένοις ὀρθογωνίοις.



Ἐστωσαν δύο εὐθεῖαι αἱ Α, ΒΓ, καὶ τετμήσθω ἡ ΒΓ, ὡς ἔτυχεν, κατὰ τὰ Δ, Ε σημεῖα· λέγω, ὅτι τὸ ὑπὸ τῶν Α, ΒΓ περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶν ὑπὸ τῶν Α, ΒΔ περιεχομένῳ ὀρθογωνίῳ καὶ τῶν ὑπὸ τῶν Α, ΔΕ καὶ ἔτι τῶν ὑπὸ τῶν Α, ΕΓ.

Ἦχθω γὰρ ἀπὸ τοῦ Β τῆς ΒΓ πρὸς ὀρθὰς ἡ ΒΖ, καὶ κείσθω τῆς Α ἴση ἡ ΒΗ, καὶ διὰ μὲν τοῦ Η τῆς ΒΓ παράλληλος ἦχθω ἡ ΗΘ, διὰ δὲ τῶν Δ, Ε, Γ τῆς ΒΗ παράλληλοι ἦχθωσαν αἱ ΔΚ, ΕΛ, ΓΘ.

ἴσον δὲ ἐστὶ τὸ ΒΘ τοῖς ΒΚ, ΔΛ, ΕΘ. καὶ ἐστὶ τὸ μὲν ΒΘ τὸ ὑπὸ τῶν Α, ΒΓ· περιέχεται μὲν γὰρ ὑπὸ τῶν ΗΒ, ΒΓ, ἴση δὲ ἡ ΒΗ τῆς Α· τὸ δὲ ΒΚ τὸ ὑπὸ τῶν Α, ΒΔ· περιέχεται μὲν γὰρ ὑπὸ τῶν ΗΒ, ΒΔ, ἴση δὲ ἡ ΒΗ τῆς Α. τὸ δὲ ΔΛ τὸ ὑπὸ τῶν Α, ΔΕ· ἴση γὰρ ἡ ΔΚ, τουτέστιν ἡ ΒΗ, τῆς Α. καὶ ἔτι ὁμοίως τὸ ΕΘ τὸ ὑπὸ τῶν Α, ΕΓ· τὸ ἄρα ὑπὸ τῶν Α, ΒΓ ἴσον ἐστὶ τῶν ὑπὸ Α, ΒΔ καὶ τῶν ὑπὸ Α, ΔΕ καὶ ἔτι τῶν ὑπὸ Α, ΕΓ.

Ἐὰν ἄρα ὡσι δύο εὐθεῖαι, τμηθῆ δὲ ἡ ἑτέρα αὐτῶν εἰς ὅσα δηποτοῦν τμήματα, τὸ περιεχόμενον ὀρθογώνιον ὑπὸ τῶν δύο εὐθειῶν ἴσον ἐστὶ τοῖς ὑπὸ τε τῆς ἀτμήτου καὶ ἐκάστου τῶν τμημάτων περιεχομένοις ὀρθογωνίοις· ὅπερ


Definitions

1. Any rectangular parallelogram is said to be contained by the two straight-lines containing the right-angle.

2. And in any parallelogrammic figure, let any one whatsoever of the parallelograms about its diagonal, (taken) with its two complements, be called a gnomon.

Proposition 1†

If there are two straight-lines, and one of them is cut into any number of pieces whatsoever, then the rectangle contained by the two straight-lines is equal to the (sum of the) rectangles contained by the uncut (straight-line), and every one of the pieces (of the cut straight-line).



Let A and BC be the two straight-lines, and let BC be cut, at random, at points D and E. I say that the rectangle contained by A and BC is equal to the rectangle(s) contained by A and BD, by A and DE, and, finally, by A and EC.

For let BF have been drawn from point B, at right-angles to BC [Prop. 1.11], and let BG be made equal to A [Prop. 1.3], and let GH have been drawn through (point) G, parallel to BC [Prop. 1.31], and let DK, EL, and CH have been drawn through (points) D, E, and C (respectively), parallel to BG [Prop. 1.31].

So the (rectangle) BH is equal to the (rectangles) BK, DL, and EH. And BH is the (rectangle contained) by A and BC. For it is contained by GB and BC, and BG (is) equal to A. And BK (is) the (rectangle contained) by A and BD. For it is contained by GB and BD, and BG (is) equal to A. And DL (is) the (rectangle contained) by A and DE. For DK, that is to say BG [Prop. 1.34], (is) equal to A. Similarly, EH (is) also the (rectangle contained) by A and EC. Thus, the (rectangle contained) by A and BC is equal to the (rectangles contained) by A

ἔδει δεῖξαι.


and BD , by A and DE , and, finally, by A and EC .

Thus, if there are two straight-lines, and one of them is cut into any number of pieces whatsoever, then the rectangle contained by the two straight-lines is equal to the (sum of the) rectangles contained by the uncut (straight-line), and every one of the pieces (of the cut straight-line). (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity: $a(b + c + d + \dots) = ab + ac + ad + \dots$.

β'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑκατέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ἀπὸ τῆς ὅλης τετραγώνῳ.



Εὐθεῖα γὰρ ἡ AB τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ Γ σημεῖον· λέγω, ὅτι τὸ ὑπὸ τῶν $AB, B\Gamma$ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ὑπὸ $BA, A\Gamma$ περιεχομένου ὀρθογώνιου ἴσον ἐστὶ τῶ ἀπὸ τῆς AB τετραγώνῳ.


Ἀναγεγράφθω γὰρ ἀπὸ τῆς AB τετράγωνον τὸ $A\Delta EB$, καὶ ἦχθω διὰ τοῦ Γ ὀποτέρᾳ τῶν $A\Delta, BE$ παράλληλος ἡ ΓZ .

Ἴσον δὴ ἐστὶ τὸ AE τοῖς $AZ, \Gamma E$. καὶ ἐστὶ τὸ μὲν AE τὸ ἀπὸ τῆς AB τετράγωνον, τὸ δὲ AZ τὸ ὑπὸ τῶν $BA, A\Gamma$ περιεχόμενον ὀρθογώνιον· περιέχεται μὲν γὰρ ὑπὸ τῶν $\Delta A, A\Gamma$, ἴση δὲ ἡ $A\Delta$ τῇ AB · τὸ δὲ ΓE τὸ ὑπὸ τῶν $AB, B\Gamma$ · ἴση γὰρ ἡ BE τῇ AB . τὸ ἄρα ὑπὸ τῶν $BA, A\Gamma$ μετὰ τοῦ ὑπὸ τῶν $AB, B\Gamma$ ἴσον ἐστὶ τῶ ἀπὸ τῆς AB τετραγώνῳ.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑκατέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ἀπὸ τῆς ὅλης τετραγώνῳ· ὅπερ ἔδει δεῖξαι.

Proposition 2†

If a straight-line is cut at random then the (sum of the) rectangle(s) contained by the whole (straight-line), and each of the pieces (of the straight-line), is equal to the square on the whole.



For let the straight-line AB have been cut, at random, at point C . I say that the rectangle contained by AB and BC , plus the rectangle contained by BA and AC , is equal to the square on AB .

For let the square $ADEB$ have been described on AB [Prop. 1.46], and let CF have been drawn through C , parallel to either of AD or BE [Prop. 1.31].


So the (square) AE is equal to the (rectangles) AF and CE . And AE is the square on AB . And AF (is) the rectangle contained by the (straight-lines) BA and AC . For it is contained by DA and AC , and AD (is) equal to AB . And CE (is) the (rectangle contained) by AB and BC . For BE (is) equal to AB . Thus, the (rectangle contained) by BA and AC , plus the (rectangle contained) by AB and BC , is equal to the square on AB .

Thus, if a straight-line is cut at random then the (sum of the) rectangle(s) contained by the whole (straight-line), and each of the pieces (of the straight-line), is equal to the square on the whole. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity: $ab + ac = a^2$ if $a = b + c$.

γ'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ τε ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογωνίῳ καὶ τῶ ἀπὸ τοῦ προειρημένου τμήματος τετραγώνῳ.




Εὐθεῖα γὰρ ἡ AB τετιμήσθω, ὡς ἔτυχεν, κατὰ τὸ Γ . λέγω, ὅτι τὸ ὑπὸ τῶν AB , $B\Gamma$ περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ τε ὑπὸ τῶν $A\Gamma$, ΓB περιεχομένῳ ὀρθογωνίῳ μετὰ τοῦ ἀπὸ τῆς $B\Gamma$ τετραγώνου.

Ἄναγεγράφθω γὰρ ἀπὸ τῆς ΓB τετράγωνον τὸ $\Gamma\Delta E B$, καὶ διήχθω ἡ $E\Delta$ ἐπὶ τὸ Z , καὶ διὰ τοῦ A ὁποτέρᾳ τῶν $\Gamma\Delta$, BE παράλληλος ἤχθω ἡ AZ . ἴσον δὲ ἐστὶ τὸ AE τοῖς $A\Delta$, ΓE : καὶ ἐστὶ τὸ μὲν AE τὸ ὑπὸ τῶν AB , $B\Gamma$ περιεχόμενον ὀρθογώνιον· περιέχεται μὲν γὰρ ὑπὸ τῶν AB , BE , ἴση δὲ ἡ BE τῇ $B\Gamma$: τὸ δὲ $A\Delta$ τὸ ὑπὸ τῶν $A\Gamma$, ΓB : ἴση γὰρ ἡ $\Delta\Gamma$ τῇ ΓB : τὸ δὲ ΔB τὸ ἀπὸ τῆς ΓB τετράγωνον: τὸ ἄρα ὑπὸ τῶν AB , $B\Gamma$ περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ὑπὸ τῶν $A\Gamma$, ΓB περιεχομένῳ ὀρθογωνίῳ μετὰ τοῦ ἀπὸ τῆς $B\Gamma$ τετραγώνου.

Ἐάν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ τε ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογωνίῳ καὶ τῶ ἀπὸ τοῦ προειρημένου τμήματος τετραγώνῳ· ὅπερ ἔδει δεῖξαι.

Proposition 3†

If a straight-line is cut at random then the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the rectangle contained by (both of) the pieces, and the square on the aforementioned piece.



For let the straight-line AB have been cut, at random, at (point) C . I say that the rectangle contained by AB and BC is equal to the rectangle contained by AC and CB , plus the square on BC .

For let the square $CDEB$ have been described on CB [Prop. 1.46], and let ED have been drawn through to F , and let AF have been drawn through A , parallel to either of CD or BE [Prop. 1.31]. So the (rectangle) AE is equal to the (rectangle) AD and the (square) CE . And AE is the rectangle contained by AB and BC . For it is contained by AB and BE , and BE (is) equal to BC . And AD (is) the (rectangle contained) by AC and CB . For DC (is) equal to CB . And DB (is) the square on CB . Thus, the rectangle contained by AB and BC is equal to the rectangle contained by AC and CB , plus the square on BC .

Thus, if a straight-line is cut at random then the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the rectangle contained by (both of) the pieces, and the square on the aforementioned piece. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity: $(a + b)a = ab + a^2$.


δ'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς ὅλης τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν τμημάτων τετραγώνοις καὶ τῶ δις ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθο-

Proposition 4†

If a straight-line is cut at random then the square on the whole (straight-line) is equal to the (sum of the) squares on the pieces (of the straight-line), and twice the

γωνίω.




Εὐθεία γὰρ γραμμὴ ἡ AB τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ Γ . λέγω, ὅτι τὸ ἀπὸ τῆς AB τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν AG , GB τετραγώνοις καὶ τῷ δις ὑπὸ τῶν AG , GB περιεχομένῳ ὀρθογωνίῳ.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς AB τετράγωνον τὸ $ADEB$, καὶ ἐπεζεύχθω ἡ BD , καὶ διὰ μὲν τοῦ Γ ὁποτέρᾳ τῶν AD , EB παράλληλος ἦχθω ἡ ΓZ , διὰ δὲ τοῦ H ὁποτέρᾳ τῶν AB , DE παράλληλος ἦχθω ἡ ΘK . καὶ ἐπεὶ παράλληλός ἐστιν ἡ ΓZ τῇ AD , καὶ εἰς αὐτὰς ἐμπίπτωκεν ἡ BD , ἡ ἐκτὸς γωνία ἢ ὑπὸ ΓHB ἴση ἐστὶ τῇ ἐντὸς καὶ ἀπεναντίον τῇ ὑπὸ $A\Delta B$. ἀλλ' ἡ ὑπὸ $A\Delta B$ τῇ ὑπὸ $AB\Delta$ ἐστὶν ἴση, ἐπεὶ καὶ πλευρὰ ἡ BA τῇ $A\Delta$ ἐστὶν ἴση· καὶ ἡ ὑπὸ ΓHB ἄρα γωνία τῇ ὑπὸ $H\Gamma B$ ἐστὶν ἴση· ὥστε καὶ πλευρὰ ἡ $B\Gamma$ πλευρᾷ τῇ ΓH ἐστὶν ἴση· ἀλλ' ἡ μὲν ΓB τῇ HK ἐστὶν ἴση. ἡ δὲ ΓH τῇ KB · καὶ ἡ HK ἄρα τῇ KB ἐστὶν ἴση· ἰσόπλευρον ἄρα ἐστὶ τὸ ΓHKB . λέγω δὴ, ὅτι καὶ ὀρθογώνιον. ἐπεὶ γὰρ παράλληλός ἐστιν ἡ ΓH τῇ BK [καὶ εἰς αὐτὰς ἐμπίπτωκεν εὐθεῖα ἡ ΓB], αἱ ἄρα ὑπὸ $K\Gamma B$, $H\Gamma B$ γωνίαί δύο ὀρθαῖς εἰσὶν ἴσαι. ὀρθὴ δὲ ἡ ὑπὸ $K\Gamma B$ · ὀρθὴ ἄρα καὶ ἡ ὑπὸ $B\Gamma H$ · ὥστε καὶ αἱ ἀπεναντίον αἱ ὑπὸ ΓHK , HKB ὀρθαῖς εἰσὶν. ὀρθογώνιον ἄρα ἐστὶ τὸ ΓHKB · ἐδείχθη δὲ καὶ ἰσόπλευρον· τετράγωνον ἄρα ἐστίν· καὶ ἐστὶν ἀπὸ τῆς ΓB . διὰ τὰ αὐτὰ δὴ καὶ τὸ ΘZ τετράγωνόν ἐστιν· καὶ ἐστὶν ἀπὸ τῆς ΘH , τουτέστιν [ἀπὸ] τῆς AG · τὰ ἄρα ΘZ , $K\Gamma$ τετράγωνα ἀπὸ τῶν AG , GB εἰσὶν. καὶ ἐπεὶ ἴσον ἐστὶ τὸ AH τῷ HE , καὶ ἐστὶ τὸ AH τὸ ὑπὸ τῶν AG , GB · ἴση γὰρ ἡ $H\Gamma$ τῇ GB · καὶ τὸ HE ἄρα ἴσον ἐστὶ τῷ ὑπὸ AG , GB · τὰ ἄρα AH , HE ἴσα ἐστὶ τῷ δις ὑπὸ τῶν AG , GB . ἐστὶ δὲ καὶ τὰ ΘZ , ΓK τετράγωνα ἀπὸ τῶν AG , GB · τὰ ἄρα τέσσαρα τὰ ΘZ , ΓK , AH , HE ἴσα ἐστὶ τοῖς τε ἀπὸ τῶν AG , GB τετραγώνοις καὶ τῷ δις ὑπὸ τῶν AG , GB περιεχομένῳ ὀρθογωνίῳ. ἀλλὰ τὰ ΘZ , ΓK , AH , HE ὅλον ἐστὶ τὸ $ADEB$, ὃ ἐστὶν ἀπὸ τῆς AB τετράγωνον· τὸ ἄρα ἀπὸ τῆς AB τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν AG , GB τετραγώνοις καὶ τῷ δις ὑπὸ τῶν AG , GB περιεχομένῳ ὀρθογωνίῳ.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς

rectangle contained by the pieces.



For let the straight-line AB have been cut, at random, at (point) C . I say that the square on AB is equal to the (sum of the) squares on AC and CB , and twice the rectangle contained by AC and CB .

For let the square $ADEB$ have been described on AB [Prop. 1.46], and let BD have been joined, and let CF have been drawn through C , parallel to either of AD or EB [Prop. 1.31], and let HK have been drawn through G , parallel to either of AB or DE [Prop. 1.31]. And since CF is parallel to AD , and BD has fallen across them, the external angle CGB is equal to the internal and opposite (angle) ADB [Prop. 1.29]. But, ADB is equal to ABD , since the side BA is also equal to AD [Prop. 1.5]. Thus, angle CGB is also equal to GBC . So the side BC is equal to the side CG [Prop. 1.6]. But, CB is equal to GK , and CG to KB [Prop. 1.34]. Thus, GK is also equal to KB . Thus, $CGKB$ is equilateral. So I say that (it is) also right-angled. For since CG is parallel to BK [and the straight-line CB has fallen across them], the angles KBC and GCB are thus equal to two right-angles [Prop. 1.29]. But KBC (is) a right-angle. Thus, BCG (is) also a right-angle. So the opposite (angles) CGK and GKB are also right-angles [Prop. 1.34]. Thus, $CGKB$ is right-angled. And it was also shown (to be) equilateral. Thus, it is a square. And it is on CB . So, for the same (reasons), HF is also a square. And it is on HG , that is to say [on] AC [Prop. 1.34]. Thus, the squares HF and KC are on AC and CB (respectively). And the (rectangle) AG is equal to the (rectangle) GE [Prop. 1.43]. And AG is the (rectangle contained) by AC and CB . For GC (is) equal to CB . Thus, GE is also equal to the (rectangle contained) by AC and CB . Thus, the (rectangles) AG and GE are equal to twice the (rectangle contained) by AC and CB . And HF and CK are the squares on AC and CB (respectively). Thus, the four (figures) HF , CK , AG , and GE are equal to the (sum of the) squares on

ἄλλης τετραγώνων ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν τμημάτων τετραγώνοις καὶ τῷ δις ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογώνῳ· ὅπερ ἔδει δεῖξαι.


AC and *BC*, and twice the rectangle contained by *AC* and *CB*. But, the (figures) *HF*, *CK*, *AG*, and *GE* are (equivalent to) the whole of *ADEB*, which is the square on *AB*. Thus, the square on *AB* is equal to the (sum of the) squares on *AC* and *CB*, and twice the rectangle contained by *AC* and *CB*.

Thus, if a straight-line is cut at random then the square on the whole (straight-line) is equal to the (sum of the) squares on the pieces (of the straight-line), and twice the rectangle contained by the pieces. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity: $(a + b)^2 = a^2 + b^2 + 2ab$.

ε'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ εἰς ἴσα καὶ ἄνισα, τὸ ὑπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς μεταξὺ τῶν τομῶν τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ἡμισείας τετραγώνῳ.




Εὐθεῖα γὰρ τις ἡ *AB* τετμήσθω εἰς μὲν ἴσα κατὰ τὸ *Γ*, εἰς δὲ ἄνισα κατὰ τὸ *Δ*· λέγω, ὅτι τὸ ὑπὸ τῶν *ΑΔ*, *ΔΒ* περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς *ΓΔ* τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς *ΓΒ* τετραγώνῳ.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς *ΓΒ* τετραγώνων τὸ *ΓΕΖΒ*, καὶ ἐπεζύχθω ἡ *ΒΕ*, καὶ διὰ μὲν τοῦ *Δ* ὁποτέρᾳ τῶν *ΓΕ*, *ΒΖ* παράλληλος ἦχθω ἡ *ΔΗ*, διὰ δὲ τοῦ *Θ* ὁποτέρᾳ τῶν *ΑΒ*, *ΕΖ* παράλληλος πάλιν ἦχθω ἡ *ΚΜ*, καὶ πάλιν διὰ τοῦ *Α* ὁποτέρᾳ τῶν *ΓΛ*, *ΒΜ* παράλληλος ἦχθω ἡ *ΑΚ*. καὶ ἐπεὶ ἴσον ἐστὶ τὸ *ΓΘ* παραπλήρωμα τῷ *ΘΖ* παραπλήρωματι, κοινὸν προσκείσθω τὸ *ΔΜ*· ὅλον ἄρα τὸ *ΓΜ* ὅλω τῷ *ΔΖ* ἴσον ἐστίν. ἀλλὰ τὸ *ΓΜ* τῷ *ΑΛ* ἴσον ἐστίν, ἐπεὶ καὶ ἡ *ΑΓ* τῆ *ΓΒ* ἐστίν ἴση· καὶ τὸ *ΑΛ* ἄρα τῷ *ΔΖ* ἴσον ἐστίν. κοινὸν προσκείσθω τὸ *ΓΘ*· ὅλον ἄρα τὸ *ΑΘ* τῷ *ΜΝΞ*† γνώμονι ἴσον ἐστίν. ἀλλὰ τὸ *ΑΘ* τὸ ὑπὸ τῶν *ΑΔ*, *ΔΒ* ἐστίν ἴση γὰρ ἡ *ΔΘ* τῆ *ΔΒ*· καὶ ὁ *ΜΝΞ* ἄρα γνώμων ἴσος ἐστὶ τῷ ὑπὸ *ΑΔ*, *ΔΒ*. κοινὸν προσκείσθω τὸ *ΛΗ*, ὅ ἐστιν ἴσον τῷ ἀπὸ τῆς *ΓΔ*· ὁ ἄρα *ΜΝΞ* γνώμων καὶ τὸ *ΛΗ* ἴσα ἐστὶ τῷ ὑπὸ τῶν *ΑΔ*, *ΔΒ* περιεχομένῳ ὀρθογώνῳ καὶ τῷ ἀπὸ τῆς

Proposition 5†

If a straight-line is cut into equal and unequal (pieces) then the rectangle contained by the unequal pieces of the whole (straight-line), plus the square on the (difference) between the (equal and unequal) pieces, is equal to the square on half (of the straight-line).



For let any straight-line *AB* have been cut—equally at *C*, and unequally at *D*. I say that the rectangle contained by *AD* and *DB*, plus the square on *CD*, is equal to the square on *CB*.

For let the square *CEFB* have been described on *CB* [Prop. 1.46], and let *BE* have been joined, and let *DG* have been drawn through *D*, parallel to either of *CE* or *BF* [Prop. 1.31], and again let *KM* have been drawn through *H*, parallel to either of *AB* or *EF* [Prop. 1.31], and again let *AK* have been drawn through *A*, parallel to either of *CL* or *BM* [Prop. 1.31]. And since the complement *CH* is equal to the complement *HF* [Prop. 1.43], let the (square) *DM* have been added to both. Thus, the whole (rectangle) *CM* is equal to the whole (rectangle) *DF*. But, (rectangle) *CM* is equal to (rectangle) *AL*, since *AC* is also equal to *CB* [Prop. 1.36]. Thus, (rectangle) *AL* is also equal to (rectangle) *DF*. Let (rectangle) *CH* have been added to both. Thus, the whole (rectangle) *AH* is equal to the gnomon *NOP*. But, *AH*

ΓΔ τετραγώνω. ἀλλὰ ὁ ΜΝΞ γνώμων καὶ τὸ ΛΗ ὄλον ἐστὶ τὸ ΓΕΖΒ τετράγωνον, ὃ ἐστὶν ἀπὸ τῆς ΓΒ· τὸ ἄρα ὑπὸ τῶν ΑΔ, ΔΒ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΓΔ τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ΓΒ τετραγώνω.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ εἰς ἴσα καὶ ἄνισα, τὸ ὑπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς μεταξὺ τῶν τομῶν τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ἡμισείας τετραγώνω. ὅπερ ἔδει δεῖξαι.

is the (rectangle contained) by AD and DB . For DH (is) equal to DB . Thus, the gnomon NOP is also equal to the (rectangle contained) by AD and DB . Let LG , which is equal to the (square) on CD , have been added to both. Thus, the gnomon NOP and the (square) LG are equal to the rectangle contained by AD and DB , and the square on CD . But, the gnomon NOP and the (square) LG is (equivalent to) the whole square $CEFB$, which is on CB . Thus, the rectangle contained by AD and DB , plus the square on CD , is equal to the square on CB .

Thus, if a straight-line is cut into equal and unequal (pieces) then the rectangle contained by the unequal pieces of the whole (straight-line), plus the square on the (difference) between the (equal and unequal) pieces, is equal to the square on half (of the straight-line). (Which is) the very thing it was required to show.

† Note the (presumably mistaken) double use of the label M in the Greek text.



‡ This proposition is a geometric version of the algebraic identity: $ab + [(a + b)/2 - b]^2 = [(a + b)/2]^2$.

ζ'.

Proposition 6†

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ δίχα, προστεθῆ δέ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας, τὸ ὑπὸ τῆς ὅλης σὺν τῇ προσκειμένη καὶ τῆς προσκειμένης περιεχόμενον ὀρθόγώνιον μετὰ τοῦ ἀπὸ τῆς ἡμισείας τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς συγκειμένης ἕκ τε τῆς ἡμισείας καὶ τῆς προσκειμένης τετραγώνω.

If a straight-line is cut in half, and any straight-line added to it straight-on, then the rectangle contained by the whole (straight-line) with the (straight-line) having being added, and the (straight-line) having being added, plus the square on half (of the original straight-line), is equal to the square on the sum of half (of the original straight-line) and the (straight-line) having been added.



Εὐθεῖα γάρ τις ἡ AB τετμήσθω δίχα κατὰ τὸ Γ σημεῖον, προσκείσθω δέ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας ἡ BD . λέγω, ὅτι τὸ ὑπὸ τῶν AD , DB περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς GB τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς GD τετραγώνω.

For let any straight-line AB have been cut in half at point C , and let any straight-line BD have been added to it straight-on. I say that the rectangle contained by AD and DB , plus the square on CB , is equal to the square on CD .

Ἀναγεγράφθω γὰρ ἀπὸ τῆς GD τετράγωνον τὸ $GEZ\Delta$, καὶ ἐπεζεύχθω ἡ DE , καὶ διὰ μὲν τοῦ B σημείου ὁποτέρᾳ τῶν EG , ΔZ παράλληλος ἦχθω ἡ BH , διὰ δὲ τοῦ Θ σημείου ὁποτέρᾳ τῶν AB , EZ παράλληλος ἦχθω ἡ KM , καὶ ἔτι διὰ τοῦ A ὁποτέρᾳ τῶν GL , ΔM παράλληλος ἦχθω ἡ AK .

For let the square $CEFD$ have been described on CD [Prop. 1.46], and let DE have been joined, and let BG have been drawn through point B , parallel to either of EC or DF [Prop. 1.31], and let KM have been drawn through point H , parallel to either of AB or EF [Prop. 1.31], and finally let AK have been drawn

Ἐπεὶ οὖν ἴση ἐστὶν ἡ AG τῇ GB , ἴσον ἐστὶ καὶ τὸ AL

