

## 4/11 The Cantor Set and Function

Recall we saw that if  $f$  is differentiable on  $(a,b)$  and  $f'(c) = 0$  for all  $c \in (a,b)$ , then  $f$  is constant.

Question: Can we "relax" the hypotheses here?

Answer: Yes to some extent, but not as far as you might think! (There are some truly strange functions out there!)

In particular, today, we want to construct a famous example of a function  $f: [0,1] \rightarrow \mathbb{R}$  such that:

- (1)  $f$  is continuous on  $[0,1]$
- (2)  $f$  is differentiable "almost everywhere" on  $[0,1]$ , with  $f'(c) = 0$  whenever  $f'(c)$  exists.
- (3)  $f(0) = 0$ ,  $f(1) = 1$  (is non constant!)

Will come back to this later

The construction depends on another famous example of a subset of  $\mathbb{R}$  with unexpected properties, called the Cantor set. Here's the Cantor set.

Georg Cantor

- Start with  $C_0 = [0,1]$
- $C_1 = [0, \frac{1}{3}] \cup [\frac{2}{3}, 1]$  (i.e. remove  $(\frac{1}{3}, \frac{2}{3})$  from  $C_0$ )
- $C_2 = [0, \frac{1}{9}] \cup [\frac{2}{9}, \frac{1}{3}] \cup [\frac{2}{3}, \frac{7}{9}] \cup [\frac{8}{9}, 1]$  (i.e. ...)
- $\vdots$

def  $C$ , the Cantor set is

$$C = \bigcap_{n=1}^{\infty} C_n.$$

Observations:

(1)  $C \neq \emptyset$ , since for instance  $\frac{1}{3}, \frac{2}{3}, \frac{1}{9}, \frac{2}{9}, \dots$   
are in  $C$ .

geom. series

$$(2) \quad \frac{1}{3} + 2 \cdot \frac{1}{9} + 4 \cdot \frac{1}{27} + \dots = \sum_{u=0}^{\infty} \frac{1}{3} \cdot \left(\frac{2}{3}\right)^u$$
$$= \frac{\frac{1}{3}}{1 - \frac{2}{3}} = 1$$

So: The total length of the <sup>open</sup> intervals removed from  $[0, 1]$  to get  $C$  is 1 (!)

(3)  $C$  is "as far as possible from empty"  
In fact,  $C$  is uncountably infinite.

We can represent each real in  $[0, 1]$  as a ternary (base 3) fraction, eg.

$$\frac{1}{3} = (\cdot 1)_3 = (\cdot 0\bar{2})_3$$

$$\frac{2}{3} = (\cdot 2)_3 = (\cdot 2\bar{0})_3$$

$$\frac{1}{2} = (\cdot \bar{1})_3$$

$$\frac{1}{3} + \frac{1}{9} + \frac{1}{27} + \dots$$

$$\frac{\frac{1}{3}}{1 - \frac{1}{3}} = \frac{1}{2} \checkmark$$

$$1 = (\cdot \bar{2})_3$$

$$\frac{2}{3} + \frac{2}{9} + \frac{2}{27} + \dots$$

$$= \frac{\frac{2}{3}}{1 - \frac{1}{3}} = 1 \checkmark$$

$$\begin{array}{r} .1 \\ 2 \overline{) 1.0} \\ \underline{2} \\ 1 \end{array}$$

all  $x$  here:  
 $x = (.1 \dots)_3$

Note:  $[0, \frac{1}{9}] (\frac{1}{9}, \frac{2}{9}) [\frac{2}{9}, \frac{1}{3}] (\frac{1}{3}, \frac{2}{3}) [\frac{2}{3}, \frac{7}{9}] (\frac{7}{9}, \frac{8}{9}) [\frac{8}{9}, 1]$

all  $x$  here here  
 $\{x = (.01 \dots)_3\}$

all  $x$  here  
 $x = .21 \dots$

etc. so

$$C = \left\{ x \in [0, 1] \mid x \text{ has a ternary expansion using only the digits } 0, 2 \right\}$$

$$(4) \quad \exists \quad \begin{matrix} [0, 1] \\ \downarrow \\ f: C \end{matrix} \longrightarrow [0, 1] \quad \underline{\text{onto}}$$

$$.202220 \dots \mapsto (.101110 \dots)_2$$

$\Rightarrow$  also exists  $\bar{f}: C \rightarrow [0, 1]$  1-1 onto,  
 and hence  $C$  is uncountably infinite.

### Cantor function

$$f(x) = \begin{cases} \sum_{n=1}^{\infty} \frac{a_n/2}{2^n} & \text{if } x \in C \end{cases}$$

if  $x = (.a_1 a_2 \dots)_3$

$$\sum_{n=1}^{k-1} \frac{a_n/2}{2^n} + \frac{1}{2^k}$$

if  $x \notin C$

$.a_1 a_2 \dots 1 \dots$

so eg.  $f(x) = \frac{1}{2}$  all  $x \in (\frac{1}{3}, \frac{2}{3})$

$f(x) = \frac{1}{4}$  all  $x \in (\frac{1}{9}, \frac{2}{9})$

$f(x) = \frac{3}{4}$  all  $x \in (\frac{7}{9}, \frac{8}{9})$

↑  
 1st 1  
 in each  
 digit

$\Rightarrow f$  is constant (hence diff'able, with derivative zero) on each of the intervals removed from  $[0,1]$  to get  $C$ .

"Devil's Stair case"

continuity on  $[0,1]$ .