Mathematics 242 – Principles of Analysis Problem Set 5

Due: March 14, 2014

A' Section

- 1. For each of the following sequences, determine three different subsequences, each converging to a different limit. For each one, express your three subsequences as x_{n_k} for a suitably chosen (strictly increasing) index sequence n_k , and give an explicit formula for n_k as a function of k:
- (a) $x_n = \sin\left(\frac{n\pi}{2}\right)$ (b) $x_n = \frac{n}{5} \left[\frac{n}{5}\right]$ (as usual, [] denotes the greatest integer function)
- 2. Let $x_n = \sqrt{n}$. For each of the following sequences, either express that sequence as a subsequence of the sequence x_n for some explicit (strictly increasing) index sequence n_k , or say why that is impossible:
- (a) $\{2, 3, 4, 5, \ldots\}$ (b) $\{\sqrt{3}, \sqrt{6}, \sqrt{9}, \sqrt{12}, \ldots\}$
- (c) $\{1, 2, 4, 8, 16, 32, \ldots\}$
- 3. Let $x_n = \sin\left(\frac{n\pi}{4}\right)$ and $y_n = \cos\left(\frac{n\pi}{4}\right)$.
- (a) Find a (strictly increasing) index sequence n_k such that both x_{n_k} and y_{n_k} converge.
- (b) Find a second (strictly increasing) index sequence n_k such that both x_{n_k} and y_{n_k} diverge.
- (c) Find a third (strictly increasing) index sequence n_k such that one of x_{n_k} and y_{n_k} converges and the other diverges.

'B' Section

- 1. (True or False) If the statement is true give a proof; if it is false give a counterexample.
- (a) If x_n is a sequence of strictly positive numbers converging to 0, then x_n has a strictly decreasing subsequence x_{n_k} .
- (b) If $x_n \to 0$, then x_n contains a strictly increasing subsequence or a strictly decreasing subsequence (or both).
- (c) If x_n is a monotone increasing sequence with a bounded subsequence x_{n_k} , then x_n converges.
- 2. Consider the sequence $x_n = \cos(n)$ (where we think of n as an angle expressed in radians).
- (a) Prove that x_n has a convergent subsequence.

(b) In this part of the question we will show that x_n is not convergent, though. Suppose $\lim_{n\to\infty}\cos(n)=a$ for some real number a. Using a trig identity for $\cos(n+1)$ and considering $\lim_{n\to\infty}(\cos(n+1)-\cos(n))$, show that

$$\frac{a(\cos(1)-1)}{\sin(1)} = \lim_{n \to \infty} \sin(n).$$

But then use the sequence $\lim_{n\to\infty} (\sin(n+1) - \sin(n))$ to deduce that a=0, so $\lim_{n\to\infty} \cos(n) = \lim_{n\to\infty} \sin(n) = 0$. But this is a contradiction. Explain why to conclude the proof.

3. A cluster point of a sequence x_n is a limit of a convergent subsequence x_{n_k} . (See question 1 on the A section for examples of sequences with several different cluster points.) Show that if a_m is a convergent sequence of cluster points of a given sequence x_n , then $a = \lim_{m \to \infty} a_m$ is also a cluster point of the x_n sequence.