

MATH 136 – Calculus 2  
Practice Day on Areas Between Curves and Average Values  
March 9, 2020

*Background*

We will now return to Chapter 6 and look at some applications of integration. We will start with two rather simple first ways we can use integrals to compute other quantities.

- *Areas between curves:* If  $y = f(x)$  and  $y = g(x)$  are two graphs, with  $f(x) \geq g(x)$  for all  $x$  in  $[a, b]$  then the area between the two graphs is computed by the integral

$$\text{Area} = \int_a^b \text{top curve} - \text{bottom curve} \, dx = \int_a^b f(x) - g(x) \, dx.$$

If the graphs cross for  $x$  in the interval, then the top and bottom curve will interchange. In this case, the area can be computed by the integral

$$\text{Area} = \int_a^b |f(x) - g(x)| \, dx.$$

In practice, though, to compute areas in cases like this, we would use the interval union property and split this into two or more integrals over intervals where the top and bottom curves do not change.

- *Average values:* If  $f(x)$  is integrable on  $[a, b]$ , then we can find a number  $\bar{y}$  such that

$$\int_a^b f(x) \, dx = \bar{y}(b - a).$$

This  $\bar{y}$  is called the average value of  $f$  on the interval  $[a, b]$ . Note that we can derive the following formula for  $\bar{y}$  by dividing through by the constant  $b - a$ :

$$\bar{y} = \frac{1}{b - a} \int_a^b f(x) \, dx.$$

In other words,  $\bar{y}$  is the height of a rectangle on the same base  $[a, b]$  whose (signed) area is the same as the area of the region between the graph  $y = f(x)$  and the  $x$ -axis. If  $f(x)$  is continuous on  $[a, b]$ , then the *Mean Value Theorem for Integrals* says that  $\bar{y} = f(c)$  for some  $c$  in  $[a, b]$ .

*Questions*

1. Find the area between  $y = \sin(x)$  and  $y = \cos(x)$  for  $x$  in  $[-\pi/4, \pi/4]$ . (Draw the region to start.)
2. Find the area bounded by  $y = x^2$  and  $y = x + 2$ . (Note: This means the area of the bounded region above the parabola and below the line, but not any other area between those curves.)
3. Find the area between  $y = x^2$  and  $y = x + 2$  for  $x$  in  $[-3, 3]$ . (Draw the region.)
4. Find the average value of  $f(x) = xe^{-x}$  for  $x$  in  $[0, 3]$ . Approximate the  $x = c$  such that  $f(c) = \bar{y}$  as in the Mean Value Theorem for Integrals. (Note: There is no convenient way to solve for  $x$  in an equation  $xe^{-x} = k$  using just algebra.)
5. Find the average value of  $f(x) = \sqrt{16 - x^2}$  for  $x$  in  $[0, 3]$ . Find the  $x = c$  such that  $f(c) = \bar{y}$  as in the Mean Value Theorem for Integrals.