

MATH 136 – Calculus 2

Arc Length

March 13, 2020

Background We have now seen why the arc length of a graph $y = f(x)$ for $x = a$ to $x = b$ can be computed by the integral

$$\text{Arc Length} = \int_a^b \sqrt{1 + (f'(x))^2} dx$$

It is actually relatively rare that these integrals can be computed exactly using the FTC. Many of them require numerical approximations because the complicated integrand tends to have no elementary antiderivative. The following cases are a few of the ones that *can be* evaluated by the FTC, either because the $\sqrt{1 + (f'(x))^2}$ is rather simple, or because the $1 + (f'(x))^2$ is set up to be a perfect square(!)

Questions

1. Find the arc length of the curve $y = 4 + x^{3/2}$ for $x = 0$ to $x = 4$.
2. Find the arc length of the curve $y = \frac{x^{3/2}}{3} - x^{1/2}$ for $x = 2$ to $x = 8$.
3. The most important example of this type from a practical point of view is probably the following. Chains or wires hanging under their own weight suspended from their endpoints have the shape of a curve called the *catenary*. Up to scaling, a catenary can always be given by an function of the form

$$f(x) = \frac{e^x + e^{-x}}{2}.$$

This function is also called the *hyperbolic cosine* and there are several sections in our textbook dealing with these and related functions. Find the arc length of the catenary curve above for $x = -1$ to $x = 1$.